

# Isolation and Identification of Lactic Acid Bacteria from Fermented Soybean Pulp and Its Application as Starter Culture for Soy Yogurt

Charity Joy L. Sumagaysay<sup>1\*</sup>, James B. Sumagaysay<sup>1</sup>, Jeany Mae H. Menisterio<sup>2</sup>, Lalaine Grace M. Robles<sup>3</sup>, Reggie Y. Dela Cruz<sup>3</sup>, and Zeus S. Elumba<sup>3</sup>

<sup>1</sup>Mountain View College, College Heights, Mt. Nebo, Valencia City, Bukidnon, Philippines

<sup>2</sup>Research and Development, East-West Seed Company Incorporated, Buena Vista Farm, Zone 5, Sankan Manolo Fortich, Bukidnon, Philippines

<sup>3</sup>Institute of Biological Sciences, Central Mindanao University, Maramag, Bukidnon, Philippines

DOI: <https://doi.org/10.51583/IJLTEMAS.2025.1407000107>

Received: 17 July 2025; Accepted: 24 July 2025; Published: 14 August 2025

Food waste processing holds great potential in mitigating global food insecurity. Valorization of food waste is a sustainable practice that also lessens the effects of climate change. Isolation, identification, and characterization of lactic acid bacteria from fermented soya pulp was conducted. The isolated lactic acid bacteria species was utilized as starter culture to produce soy yogurt. This study was performed to identify LAB isolated from fermented soya pulp using 16s rRNA gene sequencing; to ferment soymilk using the LAB isolated from fermented soya pulp and develop soy yogurt; to determine the proximate nutritional composition of the soy yogurt through proximate analysis; and to evaluate the sensory characteristics of the yogurt such as color, aroma, texture, taste, and overall acceptability. The 16s rRNA gene sequencing procedure identified the isolated lactic acid bacteria species as *Pediococcus pentosaceus* Mees, 1934. Four yogurt samples were formulated using *P. pentosaceus* and a commercial starter culture. Proximate compositions such as titratable acidity, ash content, crude protein, and crude fat were determined. The soy-based yogurt drinks were similarly high in protein compared with the cow's milk counterpart. Notably, formulation 1 with the *P. pentosaceus* as the sole starter culture scored highest in crude protein. Significant difference in the average ash content between the four samples was observed. Post-hoc analyses using the Bonferroni correction revealed that formulation 1 with the isolated starter culture, *P. pentosaceus*, had the highest ash content. Sensory evaluation revealed that formulation 2 with both the isolated starter culture and commercial starter gained peak in color, taste, and overall acceptability. There was significant difference in the average aroma and texture scores between the four samples, favoring the cow's milk yogurt sample for the aroma and the soymilk yogurt with both *P. pentosaceus* and commercial starter culture for the texture. This study showed the potential of *P. pentosaceus* as a sole starter culture and in combination with commercial starter culture in producing functional foods from food waste such as soymilk-based yogurt. Refinement and standardization of the yogurt production protocol is recommended to produce the yogurt with better texture and taste. By so doing, product development from food waste can support sustainable food production and mitigate global food insecurity and climate change.

**Keywords:** isolation, identification, lactic acid bacteria, starter culture, soymilk yogurt, okara

## I. Introduction

Global food insecurity is on the rise these days. According to the Food and Agriculture Organization (FAO), around 1.3 billion tons of food made for human consumption is wasted annually. This number is equivalent to more than one-third of the total food produced worldwide. These deficits would result to wastage of resources (FAO, 2019). One solution to this pressing problem is valorization of food waste. Food waste processing methods include isolating, characterizing, and identifying lactic acid bacteria to produce fermented functional food products.

Meanwhile, most of the soybean pulp products are outcomes of fermentation using various microorganisms. Fermented foods have better shelf life, bioactive molecules, vitamins, and enhanced availability of necessary contents (Rezaca et al., 2018). Some also have therapeutic properties such as improving gut microbiota and gastrointestinal health and overcoming diabetes and cardiovascular diseases (Melini et al., 2019). In a study in 2024, the soya pulp was used as a culture medium for lactic acid bacteria to minimize improper disposal of soymilk by-product and to explore soya pulp potential as a culture medium to grow lactic acid bacteria and save the cost of buying the standard MRS medium (Retnowati et al., 2024).

Lactic acid bacteria have been reported to possess several health benefits such as improving gut health, modulating appetite, and managing weight, modulating the immune system, decreasing the severity of infection, showing antimicrobial effects on food-borne pathogens, lowering blood pressure, high radical scavenging activity, and improving cognitive health (Mathur et al., 2020). Because they are generally recognized as safe (GRAS), they are considered as one of the most important groups of microorganisms being utilized as starter cultures in producing functional food, exerting probiotic effects, and exhibiting food preservation potential (Frick et al., 2007). Due to their fermentative nature, they are used to produce dairy, meat, and vegetables fermented products (De Vuyst and Leroy, 2007). Lactic acid bacteria isolates can be identified from fermented foods which are being studied for the potential of

antimicrobial activity against foodborne pathogens and non-resistance to antibiotics (Banik et al., 2023). By far, dairy, and fermented foods provide the main source of probiotics. But because of lactose intolerance and high cholesterol levels, non-dairy fermented products are now gaining more interests (Liu et al., 2022). Lactic acid bacteria can be used against food-borne pathogens using fermentates. Food-grade cultures and undefined fermentates of lactic acid bacteria are more consumer-friendly and do not require extensive labelling, in contrast to purified fermentates which require extensive labelling as additives (Field et al., 2015).

At present, the usefulness of 16S rRNA gene sequence is expanded to food processing application where molecular characterization has recently become essential in classifying and identifying lactic acid bacteria (Ayivi and Ibrahim, 2022). This modern step of molecular identification for lactic acid bacteria isolated from soya milk or soya pulp can be done through 16S rRNA sequencing (Retnowati et al., 2024).

Several of studies have been published about yogurt production using lactic acid bacteria and cow's milk. However, in soymilk yogurt production, only very few research is available. Limited reviews mention only a few isolated lactic acid bacteria from fermented soybean pulp; mostly are from different fermented sources (Qi et al., 2021; Jiang et al., 2021). Of the few isolated lactic acid bacteria from fermented soybean pulp, *P. pentosaceus* was only used as a mixed starter culture for soymilk yogurt along with other lactic acid bacteria species. There are no published studies on using *P. pentosaceus* isolated from fermented soybean pulp as a pure starter culture for soymilk yogurt production. Hence, this study was conceived.

## **II. Materials and Methods**

### **Place and Duration of Study**

The molecular characterization of lactic acid bacteria samples was done by Macrogen Inc., 1007, 254 Beotkkot-ro, Geuncheon-gu, Seoul 08511, Rep. of Korea. The preparation of soy yogurt was conducted at the Isolation Room of the Department of Biology in Mountain View College of Seventh-day Adventists, Inc., College Heights, Mt. Nebo, Valencia City. Lyophilization or freeze-drying of the samples and proximate analysis were conducted at the Natural Science Research Institute (NSRI) and Center for Food Research, Innovation, and Extension (CFRIE) of Central Mindanao University, University Town, Musuan, Maramag, Bukidnon, Philippines. Sensory evaluation was performed by selected participants. The study started on February 5, 2025 until May 2025.

### **Molecular Characterization of LAB Isolated from Fermented Soya Pulp**

Pure lactic acid bacteria that were previously isolated from fermented soya pulp and stored in 85% MRS broth and 15% Glycerol were sent to Kinovett Scientific Solutions, Co. Then, online request for services to Macrogen Inc. was made following the procedures prescribed by Kinovett Scientific Solutions, Co. The service requested was identification, specifically 16S rRNA gene sequencing. Kinovett Scientific Solutions, Co. facilitated the air mail of the samples to Macrogen Inc., a South Korean public biotechnology company. Macrogen communicated the progress, results, and payment invoice through emails.

### **Preparation and Allocation of Starter Culture**

The method of Utami et al., (2020) was followed with some modifications. Frozen stock cultures of isolated lactic acid bacteria from fermented soya pulp was thawed and activated in 1 L 10% skim milk and 1% sucrose mixture. The mixture was incubated at 30°C for 18 hours. The 1L starter culture was allocated into three different formulations: each formulation received 100 mL of the starter culture per replicate. There were three replicates per formulation.

### **Preparation of Soy Yogurt**

For this procedure, the method of Ugwona et al., (2018) was followed with some modifications. The following set-ups were made: Control (Soymilk + Yakult), Formulation 1 (Soymilk + Starter Culture), Formulation 2 (Soymilk + Yakult + Starter Culture), Formulation 3 (Cow's milk + Starter Culture). Three replicates were made per set-up. One point five liter (1.5 L) of soymilk was used in all soymilk set-ups. The Cow's milk was prepared by mixing 125 g of full cream powdered milk (Bear Brand) with 1 L of distilled water. The liquid Cow's milk and soymilk samples were separately pasteurized at 85°C for 15 mins and cooled at 44°C. After pasteurization, formulations 1-3 were inoculated with 100 mL (6.7%) starter culture per replicate at 44°C. Four (4) bottles of Yakult (80 mL per bottle) with live *Lacticaseibacillus paracasei* strain Shirota was used for each liter of soymilk in the control sample and formulation 2 sample. One-hundred milliliters (100 mL) of starter culture was mixed with soymilk in formulation 1 and another 100 mL of starter culture was added into the soymilk and Yakult mixture in formulation 2. One-hundred milliliters (100 mL) of starter culture was also mixed with Cow's milk in formulation 3. All the samples were fermented in a water bath at 44°C for 7h and then allowed to cool gradually.

### **Allocation of Samples for Proximate Analysis and Nutrition Facts Analysis**

Fresh soy yogurt samples were brought to the Center for Food Research, Innovation, and Extension (CFRIE) of Central Mindanao University for immediate analysis of titratable acidity and ash content. Freeze-drying at -86 °C for 42 hours was performed using the lyophilizer at the Tuklas Lunas Development Center (TLDC). Freeze drying was done for the analysis of percentage crude fat and crude protein.

## Sensory Evaluation

Ten (10) semi-trained panelists performed the sensory evaluation assessing the attributes of the yogurt such as: color, aroma, texture, taste, and overall acceptability. Their feedback was recorded using a hedonic scale, ranging from 1 (extremely disliked) to 9 (extremely liked). Labels of the finish product were concealed from the panelists to prevent bias (Asghar et al., 2022).

## Statistical Analysis

Data were analyzed using Microsoft Excel (2016). Analysis of Variance (ANOVA) was used to compare means at  $p < 0.05$  level of significance.

## III. Results and Discussion

### Identification of the LAB Isolated from Fermented Soya Pulp through 16s rRNA Sequencing

Macrogen Inc. identified the lactic acid bacteria species that was isolated from the fermented soya pulp as *Pediococcus pentosaceus* Mees, 1934 (Figure 1). *Pediococcus* is a genus of Gram-positive lactic acid bacteria, previously placed within the family of Streptococcaceae (Chen et al., 2020), and recently classified under the family of Lactobacillaceae (d'Acerno et al., 2025). *P. pentosaceus* is one of the 12 species of *Pediococcus* that have been described; and along with *P. acidilactici*, are associated with dairy products such as cheese, fermentation of milk, and cheese manufacture (Holland et al., 2010). *Pediococcus* species usually occur in pairs or tetrads, and divide along two planes of symmetry. They are coccus shaped microbes, non-motile, and non-spore forming. The end product of their metabolism is lactic acid, making them acid tolerant, and thus classifying them under lactic acid bacteria. They are also responsible for the fermentation of cabbage, making it sauerkraut, together with *Leuconostoc* and *Lactobacillus* (Macrogen, 2025). Previous studies confirmed that *P. pentosaceus* occurs naturally in spontaneously fermented products that could have a considerable role in quality, safety, and efficiency when put together with other bacteria in a mixed fermenter (Gong and Qi, 2020; Montemurro et al., 2020; Xu et al., 2021). Like most lactic acid bacteria, *P. pentosaceus* are anaerobic and ferment sugars. They can be found in plant materials, ripened cheese, and a variety of processed meats (Macrogen, 2025).




Figure 1. Molecular identification of the isolated lactic acid bacteria from fermented soya pulp using 16S rRNA gene sequencing. a) Sample from moist colony. b) Sample from mucoid colony. c) Sample from round colony. d) Sample from irregularly shape colony.

*Pediococcus pentosaceus* has many health benefits such as probiotic effects, rich in antioxidants, lowering cholesterol, and boosting the immune system. Recently, *P. pentosaceus*' potential as an ingredient in probiotic fermentation products increasingly attracts more attention (Qi et al., 2021). A study in 2022 stated that *P. pentosaceus* is increasingly studied in the dairy industry due to its promising qualities as a starter culture bacterium. The study explored the physiological, survivability, and genomic properties of *P. pentosaceus*. The strain revealed amazing tolerance in low pH and high concentrations of bile salts. It also showed 71% and 84% activity against pathogens, strong biofilm formation, and notable resistance to antibiotics. The whole genome sequence also showed various genes responsible for stress tolerance, adhesion, biofilm formation, as well as acids and bile salts tolerance (Mgomi et al., 2022).

The use of *P. pentosaceus* from different sources is widespread in several applications such as antioxidant and fatty acid profiling of fermented goat, camel, and cow's milk (Balakrishnan and Agrawal, 2014); as an adjuvant therapeutic option for patients with ulcerative colitis (UC) (Bamba et al., 2018); anti-listerial activity (Jang et al., 2014); as starter culture in sourdough bread-making

(Plessas et al., 2020); as a heat stable, bacteriocin-producing and vancomycin-sensitive lactic culture from beans (Venkateshwari et al., 2010); and producing key odorants and non-volatile metabolites in broccoli juices (Xu et al., 2021). However, in soymilk yogurt production, only very few researches have been published; and those studies only mention the use of *P. pentosaceus* as a mixed starter culture for soymilk yogurt along with other lactic acid bacteria species. In the two reviews read about *P. pentosaceus*, there was no mention about it being isolated from the fermented soybean pulp, but from other different fermented sources (Qi et al., 2021; Jiang et al., 2021). There are also no published studies on using *P. pentosaceus* isolated from fermented soybean pulp as a pure starter culture for soymilk yogurt production. Hence, this study was conceived.

### Proximate Nutritional Composition of the Soy Yogurt

There were four proximate compositions determined by the Center for Food Research, Innovation, and Extension (CFRIE) of the Central Mindanao University. One was a physico-chemical component called, titratable acidity; and chemical components such as ash, crude protein, and crude fat. All were determined in percentage form (Table 1).


**Table 1. Mean percentage and standard deviation values of the proximate nutritional compositions of the soy yogurt drink samples.**

| Proximate Composition     | Control (%)            | F1 (%)                  | F2 (%)                 | F3 (%)                   |
|---------------------------|------------------------|-------------------------|------------------------|--------------------------|
| <b>Titratable Acidity</b> | 0.73±0.009             | 0.5±0.02                | 0.61±0.02              | 0.43±0.01                |
| <b>Ash Content</b>        | 5.63±0.08 <sup>a</sup> | 6.13±0.10 <sup>bd</sup> | 5.89±0.82 <sup>f</sup> | 3.37±0.24 <sup>ceg</sup> |
| <b>Crude Protein</b>      | 15.4±0.25              | 19.63±0.09              | 18.8±0.73              | 15.45±0.12               |
| <b>Crude Fat</b>          | 1.73±0.08              | 7.61±0.50               | 4.59±0.22              | 15.02±0.15               |

\*mean value of the same letter are not significantly different at P value 0.05.

The titratable acidity, crude protein, and crude fat of the samples were comparably similar. The soymilk based samples however showed high fermentation activity indicated in the titratable acidity. This meant that soymilk based samples produced more lactic acid compared to the control. Meanwhile, the soy-based yogurt drinks were similarly high in protein compared with the cow's milk counterpart. Notably, formulation 1 with the *P. pentosaceus* as the sole starter culture scored highest in crude protein. This meant that soy-based yogurts can be good protein source alternatives for cow's milk yogurt. Additionally, the results revealed that *P. pentosaceus* can be used solely as a starter culture to develop equally protein rich, plant-based alternative functional food from food waste.

There was significant difference in the average ash content between the four samples. Post-hoc analyses using the Bonferroni correction revealed that the amount of ash in the control group scored significantly lower compared with the ash of formulation 1. The ash of the control group scored significantly higher than formulation 3. The ash of formulation 1 also scored significantly higher than formulation 3; and the ash of formulation 2 also scored significantly higher than formulation 3. Notably, formulation 1 with the *P. pentosaceus* as the sole starter culture scored highest in the ash content. This may suggest higher mineral content that is naturally present in formulation 1 (Figure 2).



**Figure 2. Proximate nutritional components of the soymilk-based yogurt drink samples: control, formulation 1, and formulation 2; in comparison with formulation 3 as the cow's milk-based yogurt drink counterpart. All the results are presented as mean ± standard deviation of the mean (n=3).**

### Sensory Characteristics of the Yogurt

A 9-point hedonic scale sensory evaluation of the formulated yogurt drink samples was used. There were five sensory parameters considered: color, aroma, texture, taste, and overall acceptability (Table 2).

The color, taste, and overall acceptability of the yogurt drink samples were comparably similar. However, formulation 2 (*P. pentosaceus* with commercial starter culture) scored highest in color, taste, and overall acceptability. Although using different lactic acid bacteria species, this result is similar with Olusola and Adepoju (2022), where the soy yogurt fermented by both the isolated lactic acid bacteria starter culture and the commercial starter culture scored higher in sensory evaluation in contrast to the soy yogurt solely fermented by the isolated lactic acid bacteria.

**Table 2. Mean and standard deviation values of the proximate nutritional compositions of the soy yogurt drink samples.**

| Sensory Parameters     | Control               | F1                    | F2                    | F3                  |
|------------------------|-----------------------|-----------------------|-----------------------|---------------------|
| Color                  | 6.5±0.97              | 6.5±1.18              | 7.3±1.16              | 6.6±1.65            |
| Aroma                  | 6.9±0.74 <sup>a</sup> | 5.5±0.97 <sup>b</sup> | 6.8±1.55 <sup>a</sup> | 7±1.25 <sup>a</sup> |
| Texture                | 5.2±0.42 <sup>a</sup> | 6.1±1.29 <sup>a</sup> | 6.3±1.06 <sup>b</sup> | 5±0.82 <sup>c</sup> |
| Taste                  | 6.3±0.67              | 6±0.47                | 7.5±0.53              | 4.6±0.52            |
| Over-All Acceptability | 7.1±0.88              | 5.8±0.79              | 7.2±0.79              | 3.1±1.10            |

There was significant difference in the average aroma scores between the four samples. Post-hoc analyses using the Bonferroni correction revealed that the aroma of the control group scored significantly higher compared with the aroma of formulation 1 (Tables 3 and 4).

**Table 3. Mean and variance values of the aroma scores in the yogurt drink samples.**

| Groups  | Count | Sum | Average | Variance |
|---------|-------|-----|---------|----------|
| Control | 10    | 69  | 6.9     | 0.544444 |
| F1      | 10    | 55  | 5.5     | 0.944444 |
| F2      | 10    | 68  | 6.8     | 2.4      |
| F3      | 10    | 70  | 7       | 1.555556 |

**Table 4. ANOVA of the aroma scores in the yogurt drink samples.**

| Source of Variation | SS   | df | MS       | F       | P-value | F crit     |
|---------------------|------|----|----------|---------|---------|------------|
| Between Groups      | 14.9 | 3  | 4.966667 | 3.64898 | 0.02143 | 2.86626555 |
| Within Groups       | 49   | 36 | 1.361111 |         |         |            |
| Total               | 63.9 | 39 |          |         |         |            |

There was also significant difference in the average texture scores between the four samples. Post-hoc analyses using the Bonferroni correction revealed that the texture of the control group scored significantly lower compared with the texture of formulation 2; and the texture of formulation 2 scored significantly higher than formulation 3 (Tables 5 and 6).

**Table 5. Mean and ANOVA values of the texture scores in the yogurt drink samples.**

| Groups  | Count | Sum | Average | Variance |
|---------|-------|-----|---------|----------|
| Control | 10    | 52  | 5.2     | 0.177778 |
| T1      | 10    | 61  | 6.1     | 1.655556 |
| T2      | 10    | 63  | 6.3     | 1.122222 |
| T3      | 10    | 50  | 5       | 0.666667 |



Table 6. Mean and ANOVA values of the texture scores in the yogurt drink samples.

| Source of Variation | SS   | df | MS       | F        | P-value | F crit   |
|---------------------|------|----|----------|----------|---------|----------|
| Between Groups      | 12.5 | 3  | 4.166667 | 4.601227 | 0.00795 | 2.866266 |
| Within Groups       | 32.6 | 36 | 0.905556 |          |         |          |
| Total               | 45.1 | 39 |          |          |         |          |

Over-all, there were significant differences in the aroma and texture of the yogurt drink samples. Formulation 1 (*P. pentosaceus*) scored lowest in odor, while formulation 3 (cow's milk) revealed the most favored aroma. The lower preference in aroma for soy-based yogurt may be caused by its naturally, slightly beany or grassy smell. Meanwhile, formulation 2 (*P. pentosaceus* and commercial starter culture) scored highest in texture followed by formulation 1 (*P. pentosaceus*) (Figure 3). This result is similar with Olusola and Adepoju (2022).




Figure 3. Sensory evaluation on attributes: color, aroma, texture, taste, and over-all acceptability of soymilk-based and cow's milk-based yogurt drinks. All the results are presented as mean ± standard deviation of the mean (n=10).

#### IV. Conclusion and Recommendation

Valorization of food waste using the naturally fermented soybean pulp as the prebiotic ingredient in isolating lactic acid bacteria with a probiotic potential was conducted prior to this study. Results revealed that the isolated lactic acid bacterium was identified as *Pediococcus pentosaceus* Mees, 1934. This species was utilized as a starter culture in producing the soymilk yogurt drinks in this study. Proximate analysis showed that formulation 1 with the *P. pentosaceus* as the sole starter culture scored highest in crude protein and ash content. It also scored second highest in crude fat. There was significant difference in the average ash content between the four samples. The color, taste, and overall acceptability of the yogurt drink samples were comparably similar. But formulation 2 scored highest in color, taste, and overall acceptability. Significant differences were observed in the average aroma and textures scores.

This study showed the potential of *P. pentosaceus* as a sole starter culture and in combination with commercial starter culture in producing functional foods from food waste such as soymilk-based yogurt. Refinement of the methods and ingredients used in the process is recommended to produce the yogurt with the desired texture and taste. It is also recommended to include the determination of moisture, pH, total soluble solids for the sweetness, and further nutritional facts analysis of the product. Further, increasing the participants for the sensory analysis is also recommended to improve the product's overall acceptability for commercial production. Bile tolerance test and antimicrobial screening against foodborne pathogens and non-resistance to antibiotics are also suggested to explore more of this species characteristics and usability. By so doing, product development from food waste can support sustainable food production and mitigate global food insecurity and climate change.

#### References

1. Aritonang, Salam N., Elly Roza, Evy Rossi, Endang Purwati and Husmaini. (2017). Isolation and identification of lactic acid bacteria from okara and evaluation of their potential as candidate probiotics. Pak. J. Nutr., 16: 618-628.
2. Asghar, Aasma; Afzaal, Muhammad; Nosheen, Farhana; Saeed, Farhan; Nayik, Gulzar Ahmad; AL-Farga, Ammar; Alansari, Wafa S.; Eskandrani, Areej A.; and Shamlan, Ghalia. (2022). Isolation and Molecular Characterization of

- Processed Soybean Waste for the Development of Synbiotic Yogurt. Fermentation, 8, 622. <https://doi.org/10.3390/fermentation8110622>.
3. Ayivi, R. D., Gyawali, R., Krastanov, A., Aljaloud, S. O., Worku, M., Tahergorabi, R., Silva, R. C. d., & Ibrahim, S. A. (2020). Lactic acid bacteria: Food safety and human health applications. Dairy, 1(3), 202–232. <https://doi.org/10.3390/dairy1030015>
4. Ayivi, Raphael D. and Ibrahim, Salam A. (2022). Lactic acid bacteria: an essential probiotic and starter culture for the production of yoghurt. International Journal of Food Science and Technology, 57:7008–7025.
5. Axelsson, L. (2004). Lactic acid bacteria: classification and physiology. Food Science and Technology-New York-Marcel Dekker. 139:1-66.
6. Balakrishnan, G., and Agrawal, R. (2014). Antioxidant activity and fatty acid profile of fermented milk prepared by *Pediococcus pentosaceus*. J. Food Sci. Technol. 51, 4138–4142. doi: 10.1007/s13197-012-0891-9
7. Bamba, S., Takahashi, K., Imaeda, H., Nishida, A., Kawahara, M., Inatomi, O., et al. (2018). Effect of fermented vegetable beverage containing *Pediococcus pentosaceus* in patients with mild to moderate ulcerative colitis. Biomed. Rep. 9, 74–80. doi: 10.3892/br.2018.1099
8. Banik, A., Anjum, H., Habib, H., Abony, M., Begum, A., Ahmed, Z. (2023). Characterization of lactic acid bacteria isolated from street pickles of Dhaka, Bangladesh. Heliyon 9 e17508.
9. Bintsis, Thomas. (2018). Lactic acid bacteria as starter cultures : An update in their metabolism and genetics. AIMS Microbiology, 4(4): 665–684. DOI: 10.3934/microbiol.2018.4.665
10. Bren d'Amour, C., Reitsma, F., Baiocchi, G., Barthel, S., Güneralp, B., Erb, K.-H., . . . Seto, K. C. (2017). Future urban land expansion and implications for global croplands. Proceedings of the National Academy of Sciences, 114(34), 8939. doi:http://doi.org/10.1073/pnas.1606036114
11. Chen, Y. S., F. Yangida & T. Shinahara. (2005). Isolation and identification of lactic acid bacteria from soil using an enrichment procedure. Lett. Applied Microbiol., 40: 195-200.
12. Chen, T., Wang, L., Li, Q., Long, Y., Lin, Y., Yin, J., et al. (2020). Functional probiotics of lactic acid bacteria from Hu sheep milk. BMC Microbiol. 20:228. doi: 10.1186/s12866-020-01920-6
13. Cirlini, M., Ricci, A., Galaverna, G., & Lazzi, C. (2020). Application of lactic acid fermentation to elderberry juice: Changes in acidic and glucidic fractions. LWT, 118, 108779.
14. d'Acerno A, Nazzaro F, Reale A, Sorrentino A, Rossi M, Bucci G, Ianigro M. (2025). CNR-ISA Agri-food Microorganism Dataset (ISA01-CC). Version 1.1. Consiglio Nazionale delle Ricerche, Istituto di Bioscienze e BioRisorse (CNR-IBBR). Occurrence dataset <https://doi.org/10.15468/2ze5cv> accessed via GBIF.org on 2025-04-20. <https://gbif.org/occurrence/4920304437>
15. Daggett, P-M. and Simone, FP. (1989). Method of culturing freeze-dried microorganisms and resultant preparation. US4879239A United States.
16. Dash, Chinmaya & Payyappilli, Rajan. (2016). KOH string and Vancomycin susceptibility test as an alternative method to Gram staining. Journal of International Medicine and Dentistry. 3. 88-90. 10.18320/JIMD/201603.0288.
17. Dawe, D., Dobermann, A., Moya, P., Abdurachman, S., Singh, B., Lal, P., . . . Zhen, Q. X. (2000). How widespread are yield declines in long-term rice experiments in Asia? Field Crops Research, 66(2), 175-193. doi:https://doi.org/10.1016/S0378-4290(00)00075-7.
18. De Vuyst L & Leroy F. (2007). Bacteriocins from Lactic Acid Bacteria: Production, Purification, and Food Applications. J Mol Microbiol Biotechnol. 13: 194-199. FAO. (2019). Food Loss and Food Waste. Retrieved from <http://www.fao.org/foodloss-and-food-waste/en/>
19. Elida, M., A. Agustina, E. Ermiati, and S. Desminarte. (2022). Isolate Characterization and Amylolytic Properties of Lactic Acid Bacteria from Traditional Fermented Dadih. 1st Lekantara Annual Conference on Natural Science and Environment (LeNS 2021). IOP Conf. Series: Earth and Environmental Science 1097 (2022) 012025. IOP Publishing. doi:10.1088/1755-1315/1097/1/01202.
20. Field, D.; Daly, K.; O'Connor, P.M.; Cotter, P.D.; Hill, C.; Ross, R.P. (2015). Efficacies of Nisin A and Nisin V Semi-purified Preparations Alone and in Combination with Plant Essential Oils for Controlling *Listeria monocytogenes*. Appl. Environ. Microbiol. 2015, 81, 2762–2769.
21. Filannino, P., Di Cagno, R., & Gobbetti, M. (2018). Metabolic and functional paths of lactic acid bacteria in plant foods: Get out of the labyrinth. Current Opinion in Biotechnology, 49, 64–72.
22. Frick, Julia-Stefanie, Katrin Schenk, Matteo Quitadamo, Frauke Kahl, Martin Köberle, Erwin Bohn, Martin Aepfelbacher, & Ingo B. Autenrieth. (2007). *Lactobacillus fermentum* attenuates the proinflammatory effect of *Yersinia enterocolitica* on human epithelial cells. Inflammatory Bowel Diseases, 13(1): 83-90.
23. Farmer, S. and Alibek, K. (2019). Production and preservation of bacillus reference culture for generating standardized and reliable inocula. WO2019222168A1 WIPO (PCT).
24. Gong, Y., and Qi, X. (2020). A study revealing volatile aroma produced by *Pediococcus pentosaceus* in dough fermentation. Food Sci. Nutr. 8, 5077–5085. doi: 10.1002/fsn3.1807
25. Gobbetti, M., M. De Angelis, R. Di Cagno, L. Mancini, and P. F. Fox. (2015). Pros and cons for using non-starter lactic acid bacteria (NSLAB) as secondary/adjunct starters for cheese ripening. Trends Food Sci. Technol. 45:167–178. <https://doi.org/10.1016/j.tifs.2015.07.016>.

26. Gudisa, A. and Yenasew, A. (2022). Isolation, Identification, and Biochemical Characterization of Lactic Acid Bacteria from Okara. Food Science and Quality Management. www.iiste.org ISSN 2224-6088 (Paper) ISSN 2225-0557 (Online) Vol.117, 2022
27. Holland, R., Crow, V., & Curry, B. (2010). Lactic Acid Bacteria | *Pediococcus* spp. Encyclopedia of Dairy Sciences (Second Edition), 149-152. <https://doi.org/10.1016/B978-0-12-374407-4.00269-7>
28. Hu, Dan; Wu, Jinyong; Jin, Long; Yuan, Lixia; Li, Jun; Chen, Xiangsong; and Yao, Jianming. (2021). Evaluation of *Pediococcus pentosaceus* strains as probiotic adjunct cultures for soybean milk post-fermentation. Food Research International, Volume 148, 2021, 110570, ISSN 0963-9969, <https://doi.org/10.1016/j.foodres.2021.110570>. (<https://www.sciencedirect.com/science/article/pii/S0963996921004695>)
29. Jang, S., Lee, J., Jung, U., Choi, H. S., and Suh, H. J. (2014). Identification of an anti-listerial domain from *Pediococcus pentosaceus* T1 derived from Kimchi, a traditional fermented vegetable. Food Control 43, 42–48. doi:10.1016/j.foodcont.2014.02.040
30. Jenkins, Claire; Ling, Clare L.; Ciesielczuk, Holly L.; Lockwood, Julianne; Hopkins, Susan; McHugh, Timothy D.; Gillespie, Stephen H.; and Kibbler, Christopher C. (2012). Detection and identification of bacteria in clinical samples by 16S rRNA gene sequencing: comparison of two different approaches in clinical practice. Journal of Medical Microbiology 61(4). <https://doi.org/10.1099/jmm.0.030387-0>
31. Jiang, Shiman; Cai, Lingzhi; Lv, Longxian, and Li, Lanjuan. (2021). *Pediococcus pentosaceus*, a future additive or probiotic candidate. Microbial Cell Factories, 20:45, <https://doi.org/10.1186/s12934-021-01537-y>
32. Johnson, Jethro S.; Spakowicz, Daniel J.; Hong, Bo-Young; Petersen, Lauren M; Demkowicz, Patrick; Chen, Lei; Leopold, Shana R; Hanson, Blake M.; Agresta, Hanako O; Gerstein, Mark; Sodergren, Erica; and Weinstock, George M. (2019). Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome analysis, Nature Communications, 10:5029 | <https://doi.org/10.1038/s41467-019-13036-1> www.nature.com/naturecommunications
33. Jose, N. M., Bunt, C. R., and Hussain, M. A. (2015). Comparison of Microbiological and Probiotic Characteristics of Lactobacilli Isolates from Dairy Food Products and Animal Rumen Contents. Microorganisms, 3(2), 198-212. <https://doi.org/10.3390/microorganisms3020198>
34. Khushboo, Karnwal A and Malik, T. (2023). Characterization and selection of probiotic lactic acid bacteria from different dietary sources for development of functional foods. Front. Microbiol. 14:1170725. doi: 10.3389/fmicb.2023.1170725
35. Kwon, H.C.; Bae, H.; Seo, H.G.; Han, S.G. (2019). Chia seed extract enhances physiochemical and antioxidant properties of yogurt. J. Dairy Sci., 102, 4870–4876.
36. Leboffe, Michael J. and Burton E. Pierce. (2010). Microbiology Laboratory Theory & Application 3e. Morton Publishing Company, 772 pp.
37. Li, Mei-Na; Han, Qiang; Wang, Nan; Wang, Ting; You, Xue-Ming; Zhang, Shuai; Zhang, Cui-Cui; Shi, Yong-Qiang; Qiao, Pei-Zhuang; Man, Cheng-Lian; Feng, Teng; Li, Yue-Yue; Zhu, Zhuang; Quan, Ke-Ji; Xu, Teng-Lin; Zhang, George Fei. (2024). 16S rRNA gene sequencing for bacterial identification and infectious disease diagnosis. Biochemical and Biophysical Research Communications, Volume 739, 150974, ISSN 0006-291X. <https://doi.org/10.1016/j.bbrc.2024.150974>. (<https://www.sciencedirect.com/science/article/pii/S0006291X24015109>)
38. Liu, C., W.J. Xue, H. Ding, C. An, S.J. Ma, and Y. Liu. (2022). Probiotic potential of Lactobacillus strains isolated from fermented vegetables in Shaanxi, China, Front. Microbiol. 12, 774903, <https://doi.org/10.3389/fmicb.2021.774903>.
39. Mateos-Aparicio, I., Redondo-Cuenca, A., Villanueva-Suárez, M. J., & Zapata-Revilla, M. (2010). Pea pod, broad bean pod, and okara, potential sources of functional compounds. LWT–Food Science and Technology, 43(9), 1467–1470.
40. Mathur, H., Beresford, T.P., and Cotter, P.D. (2020). Health Benefits of Lactic Acid Bacteria (LAB) Fermentates. Nutrients 12:1679.
41. Medic, J., Atkinson, C., & Hurburgh, C. R. (2014). Current knowledge in soybean composition. Journal of the American Oil Chemists' Society, 91(3), 363–384. <https://doi.org/10.1007/s11746-013-2407-9>
42. Melini, F., Melini, V., Luziatelli, F., Ficca, A.G., and Ruzzi, M. (2019). Health-promoting components in fermented foods: an up-to-date systematic review. Nutrients 11(5):1189. <https://doi.org/10.3390/nu11051189>
43. Meanti, F.; Mussio, C.; Rocchetti, G.; Rebecchi, A.; Lucini, L.; Morelli, L. (2024). Oat Okara Fermentation: New Insights into the Microbiological and Metabolomic Characterization. Fermentation 10:545. <https://doi.org/10.3390/fermentation10110545>
44. Mgomi, Fedrick C; Yuan, Lei; Wang, Yang; Rao, Sheng-Qi; and Yang, Zhen-Quan. (2022). Physiological properties, survivability, and genomic characteristics of *Pediococcus pentosaceus* for application as a starter culture. International Journal of Dairy Technology 75 (3): 588-602. <https://doi.org/10.1111/1471-0307.12864>
45. Montemurro, M., Celano, G., De Angelis, M., Gobbetti, M., Rizzello, C. G., and Pontonio, E. (2020). Selection of non-lactobacillus strains to be used as starters for sourdough fermentation. Food Microbiol. 90:103491. doi:10.1016/j.fm.2020.103491
46. Ngouenam, R.J., Nofal, G., Patra, S., Bilkissou, N., Marius, F., Marie, K., and Francois, Zambou. (2024). Characterization of Lactic Acid Bacteria Isolated From Rotting Oranges and Use of Agropastoral Processing By-products as Carbon and Nitrogen Sources Alternative for Lactic Acid Production. BioMed Research International. 2024. 10.1155/2024/4264229.
47. Ngouénam, Romial Joel; Nofal, Ghadir; Patra, Sanjukta; Njapndounke, Bilkissou; Kouam, Edith Marius Foko; Kaktcham, Pierre Marie; and Ngoufack, François Zambou. (2024). Characterization of Lactic Acid Bacteria Isolated From Rotting



- Oranges and Use of Agropastoral Processing By-products as Carbon and Nitrogen Sources Alternative for Lactic Acid Production. Wiley BioMed Research International, Article ID 4264229, 16 pages. <https://doi.org/10.1155/2024/4264229>
48. Olusola, Abiona Stella and Adepoju, Babarinde Yinusa. (2022). The Use of Lactic Acid Bacteria (LAB) Isolates as Single and Mixed-Strain Starter Culture in Yoghurt Processing. Food Proc Nutr Sci 3(1): 15-21.
49. Othman, M., Ariff, A. B., Rios-Solis, L., & Halim, M. (2017). Extractive fermentation of lactic acid in lactic acid bacteria cultivation: A review. Frontiers in Microbiology, 8, 2285.
50. Orf, J. (2010). Introduction. In Bilyeu, K., Ratnaparkhe, M. B., & Kole, C. (Eds.), Genetics, genomics, and breeding of soybean. CRC Press.
51. Plessas, S., Mantzourani, I., and Bekatorou, A. (2020). Evaluation of *Pediococcus pentosaceus* SP2 as Starter Culture on Sourdough Bread Making. Foods 9. doi: 10.3390/foods9010077
52. Prastujati, A. U., Hilmi, M., Khusna, A., Arief, I. I., Makmur, S., and Maulida, Q. (2022). Isolation and identification of lactic acid bacteria of Bekamal (Banyuwangi traditional fermented meat). IOP Conference Series: Earth and Environmental Science, 1020, 012026. <https://doi.org/10.1088/1755-1315/1020/1/012026>.
53. Qi Y, Huang L, Zeng Y, Li W, Zhou D, Xie J, Xie J, Tu Q, Deng D and Yin J. (2021). *Pediococcus pentosaceus*: Screening and Application as Probiotics in Food Processing. Front. Microbiol. 12:762467. doi: 10.3389/fmicb.2021.762467
54. Ramadhanti, Nurazizah; Melia, Sri; Hellyward, James; and Purwati, Endang. (2021). Characteristics of lactic acid bacteria isolated from palm sugar from West Sumatra, Indonesia and their potential as a probiotic. Biodiversitas, 22(5): 2610-2616. ISSN: 1412-033X E-ISSN: 2085-4722 DOI: 10.13057/biodiv/d220520
55. Retnowati, Faizah Diah; Purwestri, Yekti Asih; and Sine, Yuni. (2024). Characterization of Lactic Acid Bacteria Isolated from Soymilk and Its Growth in Soymilk By-product Medium for the Application in Soymilk Fermentation. Journal of Tropical Biodiversity and Biotechnology, Volume 09, Issue 04: jttb89003 DOI: 10.22146/jttb.89003
56. Rezac, S., Kok, C.R., Heermann, M., and Hutkins, R. (2018). Fermented foods as a dietary source of live organisms. Front Microbiol 9: 1785, DOI: <https://doi.org/10.3389/fmicb.2018.01785>.
57. Risna, Yayuk Kurnia; Harimurti, Sri; Wihandoyo; and Widodo. (2020). Screening for probiotic of lactic acid bacteria isolated from the digestive tract of a native Aceh duck (*Anas platyrhynchos*). Biodiversitas, 21(7) : 3001-3007. ISSN: 1412-033X E-ISSN: 2085-4722 DOI: 10.13057/biodiv/d210717
58. Roslan, I.N.D.; Kamaruding, N.A.; Ismail, N.; and Shaharuddin, S. (2020). Sensory Attributes and Other Properties of Yogurt Fortified with Immobilized *Lactobacillus Plantarum* and Soybean Residue (Okara). International Journal of Probiotics and Prebiotics, Vol. 16, pp. 1–6, 2021, doi: <https://doi.org/10.37290/ijpp2641-7197.16:1-6>, ISSN 1555-1431 print; ISSN 2641-7197 online, Copyright © 2021 by New Century Health Publishers, LLC, [www.newcenturyhealthpublishers.com](http://www.newcenturyhealthpublishers.com)
59. Satari, B.; Karimi, K. (2018). Citrus processing wastes: Environmental impacts, recent advances, and future perspectives in total valorization. Resour. Conser.Recycl., 129, 153–167.
60. Schleifer, K. H., & Ludwig, W. (1995). Phylogeny of the genus *Lactobacillus* and related genera. Systematic and Applied Microbiology, 18(4): 461-467.
61. Sharma, R., P. Garg, P. Kumar, S.K. Bhatia, and S. Kulshrestha. (2020). Microbial Fermentation and Its Role in Quality Improvement of Fermented Foods. Fermentation 2020, 6(4), 106; <https://doi.org/10.3390/fermentation6040106>
62. Septembre-Malaterre, A., Remize, F., & Poucheret, P. (2018). Fruits and vegetables as a source of nutritional compounds and phytochemicals: Changes in bioactive compounds during lactic fermentation. Food Research International, 104, 86–99.
63. Sevilla, John Russel G.; Esteban, Michael Angelo S.; Iñigo, Honey Bhabes R.; Orillaza, Audrey Mae V.; and Navarro, Baby Richard R. (2021). 16S rRNA Gene Sequence Analysis of Acetic and Lactic Acid Bacteria Isolated from Philippine Sugarcane Wine (Basi). PHILIPP AGRIC SCIENTIST, 104 (1): 75-81. ISSN 0031-7454
64. Tiensin, H., Kalibata, A., & Cole, M. (2020, 08/04/2020). Ensuring Food Security in the Era of COVID-19. Retrieved from <https://www.un.org/sustainabledevelopment/blog/2020/04/ensuring-foodsecurity-covid-19/>
65. Ugwona, F.U, Obeta, N.A, Ejinkeonye, U.B and Aliyu, S.A. (2018). Production and Quality Evaluation of Soymilk Yoghurt. Nigerian Journal of Nutritional Sciences, 39 (1).
66. Utami, Tyas; Cindarbhum, Amaralida; Khuangga, Marcella C; Rahayu, Endang S; Cahyanto, Muhammad Nur; Nurfiyanti, Sri; and Zulaichah, Eni. (2020). Preparation of Indigenous Lactic Acid Bacteria Starter Cultures for Large Scale Production of Fermented Milk. Digital Press Life Sciences 2: 00010. 10th Asian Conference of Lactic Acid Bacteria. <https://doi.org/10.29037/digitalpress.22327>
67. Valin, H., Sands, R. D., van der Mensbrugghe, D., Nelson, G. C., Ahammad, H., Blanc, E., . . . Willenbockel, D. (2014). The future of food demand: understanding differences in global economic models. Agricultural Economics, 45(1), 51-67. doi:<https://doi.org/10.1111/agec.12089>
68. Venkateshwari, S., Halami, P. M., and Vijayendra, S. V. (2010). Characterisation of the heat-stable bacteriocin-producing and vancomycin-sensitive *Pediococcus pentosaceus* CFR B19 isolated from beans. Benef. Microbes 1, 159–164. doi: 10.3920/bm2009.0032
69. Wahyuni, S., Ratna, Holilah, Asranudin, and Raharjo, Raden Alip. 2016. Antioxidant Activity of Isoflavones from Tofu Pulp Waste. 6th Int'l Conference on Agriculture, Environment and Biological Sciences (ICAEB'S'16) Dec. 21-22, 2016 Kuala Lumpur (Malaysia)

70. Xu, X., Bi, S., Lao, F., Chen, F., Liao, X., and Wu, J. (2021). Comprehensive investigation on volatile and non-volatile metabolites in broccoli juices fermented by animal- and plant-derived *Pediococcus pentosaceus*. Food Chem. 341:128118. doi: 10.1016/j.foodchem.2020.128118
71. Xu, X. L. (2020). Research on key aroma identification and regulation of traditional sour curd cheese milk fan. Master's thesis. Shanghai University of Applied Technology, Shanghai, China.
72. Yamamoto, N., Shoji, M., Hoshigami, H., Watanabe, K., Watanabe, K., Takatsuzu, T., et al. (2019). Antioxidant capacity of soymilk yogurt and exopolysaccharides produced by lactic acid bacteria. Biosci. Microbiota. Food Health 38, 97–104. doi: 10.12938/bmfh.18-017
73. Yang, Yang; Xia, Yanan; Wang, Yu Rong; Sun, Li Shan; Shuang, Quan; and Zhang, Feng Mei. (2024). Optimization of lactic acid bacterial starter culture to improve the quality and flavor characteristics of traditional Hurood. J. Dairy Sci. 107:105–122. <https://doi.org/10.3168/jds.2023-23754>
74. Zhou, R.Y., X. Huang, Z. Liu, J-Y Chua, S-Q Liu. (2022). Evaluating the effect of lactic acid bacterial fermentation on salted soy whey for development of a potential novel soy sauce-like condiment. Current Research in Food Science, 5:1826-1836. Elsevier