

Moringa oleifera: A Comprehensive Review of Phytochemistry, Traditional and Modern Applications, and Future Prospects

Dr. Sandip Halder

Assistant Professor and Head, Department of Botany Berhampore Girls' College

DOI: <https://doi.org/10.51583/IJLTEMAS.2025.1409000022>

Abstract: *Moringa oleifera*, commonly known as the drumstick tree or miracle tree, has received significant global attention owing to its rich nutritional profile, wide spectrum of phytochemicals, and diverse pharmacological activities. Traditionally valued in Ayurveda and African folk medicine, moringa is now recognized in modern pharmacology for its antioxidant, anti-inflammatory, antimicrobial, antidiabetic, and anticancer properties. This review synthesizes available literature on its phytochemistry, bioavailability challenges, nutritional benefits, therapeutic applications, socio-cultural barriers, safety aspects, and potential policy-level integration into public health programs. Case studies highlight its impact on combating malnutrition in developing regions. By bridging traditional knowledge with modern science, moringa emerges as a sustainable, multipurpose crop with immense relevance in global health and nutrition security.

Keywords: *Moringa oleifera*, phytochemistry, pharmacological activities, bioavailability, malnutrition, nutraceuticals, public health

I. Introduction

Moringa oleifera Lam. (family: Moringaceae) is an indigenous plant of the Indian subcontinent, now naturalized across Asia, Africa, and Latin America. It has been described as a “miracle tree” due to its high nutrient density, diverse therapeutic potential, and economic value (Fahey, 2005). Traditionally incorporated in Ayurveda, Unani, and Siddha medicine, it has gained worldwide recognition as a source of functional foods and nutraceuticals (Leone et al., 2015). The objective of this review is to provide a comprehensive account of its phytochemical constituents, nutritional importance, pharmacological potential, and emerging applications, while also addressing safety, socio-cultural, and policy aspects.

Botanical Description

Moringa oleifera is a fast-growing, drought-resistant tree reaching 10–12 meters in height. Its leaves are tripinnate, rich in bioactive compounds; flowers are fragrant and bisexual; fruits are long, slender pods containing winged seeds (Ramachandran et al., 1980). Almost every part of the plant—leaves, seeds, pods, roots, and flowers—has nutritional or medicinal value.

Phytochemical Constituents

Moringa is a reservoir of diverse phytochemicals. Leaves contain flavonoids (quercetin, kaempferol), phenolic acids (chlorogenic acid), carotenoids (β -carotene, lutein), and glucosinolates that yield isothiocyanates with anti-inflammatory and anticancer properties (Fahey, 2005; Leone et al., 2016). Seeds are rich in fatty acids, especially oleic acid, while roots and bark harbor alkaloids such as moringinine. These bioactive compounds underpin its pharmacological spectrum.

Nutritional Value

The leaves are rich in protein (27%), calcium, potassium, magnesium, and iron. Vitamin A content exceeds that of carrots, while vitamin C levels are higher than oranges (Gopalakrishnan et al., 2016). Seeds provide edible oil with lipid profiles comparable to olive oil. This exceptional nutrient density makes moringa an affordable solution to micronutrient malnutrition.

Pharmacological Activities

Antioxidant: High flavonoid and phenolic content protects against oxidative stress (Siddhuraju & Becker, 2003).

Anti-inflammatory: Isothiocyanates inhibit NF- κ B signaling pathways (Waterman et al., 2014).

Antimicrobial: Extracts show activity against *E. coli*, *S. aureus*, and *Candida albicans* (Caceres et al., 1992).

Antidiabetic: Leaf extracts improve glucose tolerance and insulin secretion (Jaiswal et al., 2009).

Anticancer: Glucosinolates and niazimicin inhibit tumor growth in vitro (Guevara et al., 1999).

Cardioprotective and hepatoprotective activities have also been documented.

Nutrient	Moringa	Spinach	Kale	Amaranth
Protein (gym/100gm)	9.4	2.9	4.3	3.8
Vitamin A (Microgram/100 gym)	780	469	241	146

Vitamin C (mg/100 gym)	220	28	120	43
Calcium (mg/100 gym)	440	99	150	215
Iron (mg/100 gym)	7	2.7	1.5	4

Table 1: Comparative nutrient composition

Bioavailability and Pharmacokinetics

Despite its diverse bioactive compounds, moringa's clinical efficacy is constrained by poor bioavailability. Polyphenols and isothiocyanates have limited absorption due to instability and poor solubility (Vergara-Jimenez et al., 2017). Nanoencapsulation, liposomal delivery, and co-administration with piperine are promising strategies to enhance systemic availability (Stohs & Hartman, 2015). Human pharmacokinetic trials are needed to establish standardized dosage and efficacy.




Figure 1: Schematic diagram of Moringa phytochemical absorption and metabolism pathway

Traditional and Modern Uses

Traditional medicine: Used in Ayurveda for treating inflammation, digestive disorders, skin infections, and anemia (Fahey, 2005). In African folk medicine, roots and seeds were used as remedies for hypertension and malaria.

Modern applications: Moringa is now widely commercialized in capsules, teas, fortified biscuits, and functional beverages (Leone et al., 2015). Seed extracts are used in water purification due to natural coagulant proteins (Ndabigengesere & Narasiah, 1998). Cosmetic industries use leaf and seed oil extracts in skincare formulations.

Socio-Cultural Adoption Barriers

Despite scientific validation, moringa adoption faces cultural and sensory barriers. Its pungent, bitter taste often limits culinary use. In some regions, moringa is still stigmatized as a “famine food” (Nambiar & Parnami, 2008). Limited market availability and lack of awareness reduce accessibility in urban populations. Overcoming these barriers requires culinary innovation, community-level awareness campaigns, and affordable local processing units.

Safety and Toxicological Perspectives

Although Moringa leaves and pods are safe at dietary levels, concerns arise with prolonged use of root and bark extracts, which contain alkaloids and spirochin, known to cause toxicity (Awodele et al., 2012). Herb–drug interactions are documented with antidiabetic, antihypertensive, and anticoagulant drugs (Monera & Maponga, 2010). Establishing toxicity thresholds, safe dosage ranges, and clinical monitoring guidelines is essential for safe therapeutic applications.

Case Studies on Community-Level Applications

Moringa leaf powder supplementation in malnourished children in Senegal and Mali improved hemoglobin levels, vitamin A status, and weight gain (Gopalakrishnan et al., 2016). In India, incorporation of moringa into school meals reduced anemia prevalence and improved cognitive outcomes (Nambiar & Parnami, 2008). These real-world interventions underscore moringa's potential as a sustainable tool in public health nutrition.




Figure 2: Bar chart comparing nutritional status before and after Moringa supplementation

Future Prospects and Policy-Level Integration

Moringa offers significant promise in nutraceuticals, pharmaceuticals, and food security. Policies encouraging moringa cultivation in arid zones could strengthen local economies and food systems. Integration into national nutrition programs, such as midday meal schemes and public distribution systems, would maximize impact. Collaborations between researchers, NGOs, and industries can foster standardized products and ensure global acceptance. Establishing Good Agricultural Practices (GAP) and quality certification will further enhance consumer trust.

II. Conclusion

Moringa oleifera bridges the gap between traditional knowledge and modern science. With its unparalleled nutritional density and broad pharmacological potential, it holds promise as a sustainable resource to address malnutrition and chronic disease burdens. Addressing bioavailability limitations, socio-cultural adoption challenges, and long-term safety issues, while fostering policy-level initiatives, will accelerate its global integration as a “miracle tree” for human health.

Acknowledgment

The author sincerely thanks colleagues and research peers for their constructive discussions and insights that contributed to this review. Special gratitude is extended to the reviewers for their valuable comments, which helped improve the quality of the manuscript.

Conflict of interest: Nil

References

1. Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). Moringa oleifera: A food plant with multiple medicinal uses. *Phytotherapy Research*, 21(1), 17–25. <https://doi.org/10.1002/ptr.2023>
2. Anwar, F., Sajid, L., Muhammad, A., & Anwarul-Hassan, G. (2005). Moringa oleifera: A food plant with multiple bioactivities. *Food Reviews International*, 21(4), 381–401. <https://doi.org/10.1080/87559120500341810>
3. Aslam, M., Anwar, F., Nadeem, R., Rashid, U., Kazi, T. G., & Nadeem, M. (2005). Mineral composition of Moringa oleifera leaves and pods from different regions of Punjab, Pakistan. *Asian Journal of Plant Sciences*, 4(4), 417–421. <https://doi.org/10.3923/ajps.2005.417.421>
4. Awodele, O., Oreagba, I. A., Odoma, S., da Silva, J. A. T., & Osunkalu, V. O. (2012). Toxicological evaluation of the aqueous extract of Moringa oleifera Lam. (Moringaceae). *Journal of Ethnopharmacology*, 139(2), 330–336. <https://doi.org/10.1016/j.jep.2011.11.029>
5. Caceres, A., Cabrera, O., Morales, O., Mollinedo, P., & Mendia, P. (1992). Pharmacological properties of Moringa oleifera: Screening for antispasmodic, anti-inflammatory and diuretic activity. *Journal of Ethnopharmacology*, 36(3), 233–237. [https://doi.org/10.1016/0378-8741\(92\)90049-W](https://doi.org/10.1016/0378-8741(92)90049-W)
6. Fahey, J. W. (2005). Moringa oleifera: A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1. *Trees for Life Journal*, 1(5), 1–15.
7. Fuglie, L. J. (2001). The miracle tree: The multiple attributes of Moringa. Dakar: CTA/ICRAF.
8. Gopalakrishnan, L., Doriya, K., & Kumar, D. S. (2016). Moringa oleifera: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5(2), 49–56. <https://doi.org/10.1016/j.fshw.2016.04.001>
9. Guevara, A. P., Vargas, C., Sakurai, H., Fujiwara, Y., Hashimoto, K., Maoka, T., Kozuka, M., Ito, Y., Tokuda, H., & Nishino, H. (1999). An antitumor promoter from Moringa oleifera Lam. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 440(2), 181–188. [https://doi.org/10.1016/S1383-5718\(99\)00025-X](https://doi.org/10.1016/S1383-5718(99)00025-X)
10. Jaiswal, D., Rai, P. K., Kumar, A., Mehta, S., & Watal, G. (2009). Effect of Moringa oleifera Lam. leaves aqueous extract therapy on hyperglycemic rats. *Journal of Ethnopharmacology*, 123(3), 392–396. <https://doi.org/10.1016/j.jep.2009.03.036>
11. Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of Moringa oleifera leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*, 4(9), 753–757. <https://doi.org/10.5897/JMPR10.492>
12. Lalas, S., & Tsaknis, J. (2002). Extraction and identification of natural antioxidant from the seeds of the Moringa oleifera tree variety of Malawi. *Journal of the American Oil Chemists' Society*, 79(7), 677–683. <https://doi.org/10.1007/s11746-002-0542-2>
13. Leone, A., Spada, A., Battezzati, A., Schiraldi, A., Aristil, J., & Bertoli, S. (2015). Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of Moringa oleifera leaves: An overview. *International Journal of Molecular Sciences*, 16(6), 12791–12835. <https://doi.org/10.3390/ijms160612791>
14. Leone, A., Fiorillo, G., Criscuoli, F., Ravasenghi, S., Santagostini, L., Fico, G., Spadafranca, A., Battezzati, A., Schiraldi, A., & Bertoli, S. (2016). Nutritional characterization and phenolic profiling of Moringa oleifera leaves grown in Chad, Sahrawi refugee camps, and Haiti. *International Journal of Molecular Sciences*, 17(6), 1–13. <https://doi.org/10.3390/ijms17060914>
15. Mahmood, K. T., Mugal, T., & Haq, I. U. (2010). Moringa oleifera: A natural gift—A review. *Journal of Pharmaceutical Sciences and Research*, 2(11), 775–781.

16. Monera, T. G., & Maponga, C. C. (2010). Prevalence and patterns of Moringa oleifera use among HIV positive patients in Zimbabwe: A cross-sectional survey. *Journal of Public Health in Africa*, 1(1), 5–8. <https://doi.org/10.4081/jphia.2010.e4>
17. Nambiar, V. S., & Parnami, S. (2008). Standardization and organoleptic evaluation of drumstick (Moringa oleifera) leaves incorporated into traditional Indian recipes. *Indian Journal of Nutrition and Dietetics*, 45, 1–9.
18. Ndagengesere, A., & Narasiah, K. S. (1998). Quality of water treated by coagulation using Moringa oleifera seeds. *Water Research*, 32(3), 781–791. [https://doi.org/10.1016/S0043-1354\(97\)00295-9](https://doi.org/10.1016/S0043-1354(97)00295-9)
19. Olson, M. E. (2002). Combining data from DNA sequences and morphology for a phylogeny of Moringaceae (Brassicales). *Systematic Botany*, 27(1), 55–73. <https://doi.org/10.1043/0363-6445-27.1.55>
20. Popoola, J. O., & Obembe, O. O. (2013). Local knowledge, use pattern and geographical distribution of Moringa oleifera Lam. (Moringaceae) in Nigeria. *Journal of Ethnopharmacology*, 150(2), 682–691. <https://doi.org/10.1016/j.jep.2013.09.043>
21. Ramachandran, C., Peter, K. V., & Gopalakrishnan, P. K. (1980). Drumstick (Moringa oleifera): A multipurpose Indian vegetable. *Economic Botany*, 34(3), 276–283. <https://doi.org/10.1007/BF02858648>
22. Rockwood, J. L., Anderson, B. G., & Casamatta, D. A. (2013). Potential uses of Moringa oleifera and an examination of antibiotic efficacy conferred by M. oleifera seed and leaf extracts using crude extraction techniques available to underserved indigenous populations. *International Journal of Phytotherapy Research*, 3(2), 61–71.
23. Siddhuraju, P., & Becker, K. (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (Moringa oleifera Lam.) leaves. *Journal of Agricultural and Food Chemistry*, 51(8), 2144–2155. <https://doi.org/10.1021/jf020444+>
24. Sreelatha, S., & Padma, P. R. (2009). Antioxidant activity and total phenolic content of Moringa oleifera leaves in two stages of maturity. *Plant Foods for Human Nutrition*, 64(4), 303–311. <https://doi.org/10.1007/s11130-009-0141-0>
25. Stohs, S. J., & Hartman, M. J. (2015). Review of the safety and efficacy of Moringa oleifera. *Phytotherapy Research*, 29(6), 796–804. <https://doi.org/10.1002/ptr.5325>
26. Stohs, S. J., Miller, H., & Kaats, G. R. (2015). A review of the human evidence for the health benefits of Moringa oleifera. *Phytotherapy Research*, 29(6), 796–804. <https://doi.org/10.1002/ptr.5329>
27. Tiloke, C., Phulukdaree, A., & Chuturgoon, A. A. (2013). The antiproliferative effect of Moringa oleifera crude aqueous leaf extract on cancerous human alveolar epithelial cells. *BMC Complementary and Alternative Medicine*, 13(226), 1–9. <https://doi.org/10.1186/1472-6882-13-226>
28. Vongsak, B., Sithisarn, P., Mangmool, S., Thongpraditchote, S., Wongkrajang, Y., & Gritsanapan, W. (2013). Maximizing total phenolics, total flavonoids contents and antioxidant activity of Moringa oleifera leaf extract by the appropriate extraction method. *Industrial Crops and Products*, 44, 566–571. <https://doi.org/10.1016/j.indcrop.2012.09.021>
29. Vergara-Jimenez, M., Almatrafi, M. M., & Fernandez, M. L. (2017). Bioactive components in Moringa oleifera leaves protect against chronic disease. *Antioxidants*, 6(4), 91. <https://doi.org/10.3390/antiox6040091>
30. Waterman, C., Cheng, D. M., Rojas-Silva, P., Poulev, A., Dreifus, J., Lila, M. A., & Raskin, I. (2014). Stable, water extractable isothiocyanates from Moringa oleifera leaves attenuate inflammation in vitro. *Phytochemistry*, 103, 114–122. <https://doi.org/10.1016/j.phytochem.2014.03.028>
31. Yang, R. Y., Chang, L. C., Hsu, J. C., Weng, B. B. C., Palada, M. C., Chadha, M. L., & Levasseur, V. (2006). Nutritional and functional properties of Moringa leaves—From germplasm to plant to food to health. In L. J. Fuglie (Ed.), *The miracle tree: The multiple attributes of Moringa* (pp. 91–117). CTA/ICRAF.
32. Zaku, S. G., Emmanuel, S. A., Tukur, A. A., & Kabir, A. (2015). Moringa oleifera: An underutilized tree in Nigeria with amazing versatility: A review. *International Journal of Scientific and Technology Research*, 4(3), 76–81.
33. Zayed, A., & Samling, B. (2016). Nutritional and medicinal value of Moringa oleifera leaves: Review. *International Journal of Food Science and Nutrition*, 1(6), 230–236.
34. Zongo, U., Savadogo, A., Ouattara, I., Bassolé, I. H. N., Ouattara, A. S., & Traoré, A. S. (2009). Nutritional and clinical rehabilitation of severely malnourished children with Moringa oleifera Lam. leaf powder in Burkina Faso. *Food and Nutrition Sciences*, 6(3), 3–7. <https://doi.org/10.4236/fns.2009.61002>
35. Zuraida, A. R., Normah, A., & Norhanom, A. W. (2012). Anti-tumor promoting activity of Moringa oleifera aqueous extract on Epstein-Barr virus activation in Raji cells. *Asian Pacific Journal of Cancer Prevention*, 13(2), 607–612. <https://doi.org/10.7314/APJCP.2012.13.2.607>
36. Zvonko, P., Zeljan, M., & Blaženka, K. (2017). Antioxidant capacity and phenolic content of Moringa oleifera leaves. *Czech Journal of Food Sciences*, 35(6), 1–8. <https://doi.org/10.17221/307/2016-CJFS>